

Web Appendix Table 2

Prevalence, DALY and Death rates per 100,000 for total, ischemic and haemorrhagic stroke by GBD Regions in 2013

Total Stroke						
Region name	Sex	Age	DALY	95% UI	Death	95% UI
East Asia	Male	1 to 4	79.8	50.2 to 116.0	0.9	0.6 to 1.4
	Male	5 to 9	47.3	37.2 to 59.5	0.6	0.5 to 0.7
	Male	10 to 14	55.2	43.6 to 68.9	0.7	0.6 to 0.9
	Male	15 to 19	162.1	127.7 to 218.2	2.3	1.8 to 3.1
	Female	1 to 4	44.7	25.8 to 61.1	0.5	0.3 to 0.7
	Female	5 to 9	42	30.0 to 52.5	0.5	0.4 to 0.7
	Female	10 to 14	42.9	30.4 to 54.4	0.6	0.4 to 0.7
	Female	15 to 19	65.1	43.9 to 83.9	0.9	0.6 to 1.2
	Both	1 to 4	63.5	42.1 to 86.9	0.8	0.5 to 1.0
	Both	5 to 9	44.8	36.7 to 53.1	0.6	0.5 to 0.7
	Both	10 to 14	49.5	41.1 to 58.4	0.7	0.5 to 0.8
	Both	15 to 19	116.7	95.8 to 148.4	1.7	1.4 to 2.1
Southeast Asia	Male	1 to 4	58.6	39.0 to 84.0	0.7	0.5 to 1.0
	Male	5 to 9	28.3	19.5 to 40.8	0.3	0.2 to 0.5
	Male	10 to 14	46	32.9 to 63.4	0.6	0.4 to 0.8
	Male	15 to 19	162.7	121.4 to 213.2	2.3	1.7 to 3.0
	Female	1 to 4	33.4	20.5 to 48.2	0.4	0.2 to 0.6
	Female	5 to 9	30.9	22.0 to 47.3	0.4	0.3 to 0.6
	Female	10 to 14	38.6	26.7 to 63.0	0.5	0.3 to 0.8
	Female	15 to 19	76.5	54.5 to 106.0	1.1	0.8 to 1.5
	Both	1 to 4	46.3	33.2 to 62.8	0.5	0.4 to 0.7

	Both	5 to 9	29.6	22.4 to 39.8	0.4	0.3 to 0.5
	Both	10 to 14	42.4	32.5 to 58.3	0.6	0.4 to 0.8
	Both	15 to 19	120.4	97.4 to 148.6	1.7	1.4 to 2.1
Oceania	Male	1 to 4	145.5	62.5 to 310.8	1.7	0.7 to 3.7
	Male	5 to 9	105.9	60.4 to 175.8	1.3	0.8 to 2.2
	Male	10 to 14	56.2	34.5 to 86.3	0.7	0.4 to 1.1
	Male	15 to 19	230.9	147.6 to 367.8	3.3	2.1 to 5.3
	Female	1 to 4	69.1	30.7 to 142.1	0.8	0.4 to 1.7
	Female	5 to 9	47.2	25.6 to 84.5	0.6	0.3 to 1.1
	Female	10 to 14	90.2	53.2 to 147.2	1.2	0.7 to 2.0
	Female	15 to 19	48	29.8 to 77.4	0.7	0.4 to 1.1
	Both	1 to 4	108.6	51.4 to 220.2	1.3	0.6 to 2.6
	Both	5 to 9	77.6	47.9 to 124.5	1	0.6 to 1.6
	Both	10 to 14	72.6	48.9 to 106.2	1	0.6 to 1.4
	Both	15 to 19	142.3	94.1 to 216.3	2	1.3 to 3.1
Central Asia	Male	1 to 4	24.5	17.0 to 34.3	0.3	0.2 to 0.4
	Male	5 to 9	28	20.8 to 37.5	0.3	0.3 to 0.5
	Male	10 to 14	26.9	20.1 to 35.9	0.3	0.3 to 0.5
	Male	15 to 19	101.6	73.1 to 134.6	1.4	1.0 to 1.9
	Female	1 to 4	25.7	14.2 to 37.1	0.3	0.2 to 0.4
	Female	5 to 9	26.6	16.3 to 34.5	0.3	0.2 to 0.4
	Female	10 to 14	27.7	20.2 to 39.4	0.4	0.3 to 0.5
	Female	15 to 19	64.8	46.6 to 89.2	0.9	0.7 to 1.3
	Both	1 to 4	25.1	18.5 to 33.2	0.3	0.2 to 0.4
	Both	5 to 9	27.3	21.4 to 33.3	0.3	0.3 to 0.4
	Both	10 to 14	27.3	22.1 to 34.2	0.4	0.3 to 0.4
	Both	15 to 19	83.6	66.7 to 104.2	1.2	0.9 to 1.5
Central Europe	Male	1 to 4	23	14.7 to 35.9	0.3	0.2 to 0.4

	Male	5 to 9	16.5	11.6 to 22.1	0.2	0.1 to 0.3
	Male	10 to 14	26.8	20.1 to 33.8	0.3	0.2 to 0.4
	Male	15 to 19	57.4	42.4 to 72.2	0.8	0.6 to 1.0
	Female	1 to 4	21.9	13.8 to 35.4	0.3	0.2 to 0.4
	Female	5 to 9	16.9	11.6 to 22.6	0.2	0.1 to 0.3
	Female	10 to 14	26.3	18.1 to 32.6	0.3	0.2 to 0.4
	Female	15 to 19	40.3	28.4 to 49.2	0.6	0.4 to 0.7
	Both	1 to 4	22.5	15.3 to 33.6	0.3	0.2 to 0.4
	Both	5 to 9	16.7	12.5 to 21.9	0.2	0.1 to 0.3
	Both	10 to 14	26.6	20.5 to 31.6	0.3	0.3 to 0.4
	Both	15 to 19	49	38.1 to 57.5	0.7	0.5 to 0.8
Eastern Europe	Male	1 to 4	5	2.8 to 7.4	0.1	0.0 to 0.1
	Male	5 to 9	7	4.6 to 9.8	0.1	0.0 to 0.1
	Male	10 to 14	12.8	9.0 to 17.3	0.1	0.1 to 0.2
	Male	15 to 19	50.6	34.5 to 69.1	0.7	0.4 to 0.9
	Female	1 to 4	5.8	2.3 to 8.9	0.1	0.0 to 0.1
	Female	5 to 9	8.7	5.8 to 13.8	0.1	0.1 to 0.2
	Female	10 to 14	14.9	8.5 to 20.0	0.2	0.1 to 0.2
	Female	15 to 19	44.9	25.0 to 60.3	0.6	0.3 to 0.8
	Both	1 to 4	5.4	3.2 to 7.7	0.1	0.0 to 0.1
	Both	5 to 9	7.8	5.7 to 10.2	0.1	0.1 to 0.1
	Both	10 to 14	13.9	10.3 to 17.1	0.2	0.1 to 0.2
	Both	15 to 19	47.8	36.0 to 59.5	0.6	0.5 to 0.8
High-income Asia Pacific	Male	1 to 4	13.6	8.3 to 19.5	0.2	0.1 to 0.2
	Male	5 to 9	8.5	5.8 to 11.3	0.1	0.1 to 0.1
	Male	10 to 14	16.3	11.4 to 20.8	0.2	0.1 to 0.2
	Male	15 to 19	27.3	19.6 to 35.9	0.3	0.2 to 0.5
	Female	1 to 4	9.3	6.3 to 16.2	0.1	0.1 to 0.2

	Female	5 to 9	10.9	7.2 to 14.8	0.1	0.1 to 0.2
	Female	10 to 14	12.3	9.2 to 17.7	0.1	0.1 to 0.2
	Female	15 to 19	20.7	15.4 to 32.2	0.3	0.2 to 0.4
	Both	1 to 4	11.5	8.0 to 16.0	0.1	0.1 to 0.2
	Both	5 to 9	9.7	7.1 to 12.0	0.1	0.1 to 0.1
	Both	10 to 14	14.4	11.4 to 17.6	0.2	0.1 to 0.2
	Both	15 to 19	24.1	19.2 to 30.0	0.3	0.2 to 0.4
Australasia	Male	1 to 4	4.1	2.3 to 6.3	0	0.0 to 0.1
	Male	5 to 9	3.5	2.5 to 4.7	0	0.0 to 0.0
	Male	10 to 14	7.6	5.5 to 10.0	0.1	0.0 to 0.1
	Male	15 to 19	17.6	12.0 to 23.9	0.2	0.1 to 0.3
	Female	1 to 4	3.3	2.0 to 4.7	0	0.0 to 0.1
	Female	5 to 9	4	2.8 to 5.6	0	0.0 to 0.1
	Female	10 to 14	7.4	5.3 to 9.9	0.1	0.0 to 0.1
	Female	15 to 19	16.1	10.1 to 21.3	0.2	0.1 to 0.3
	Both	1 to 4	3.7	2.5 to 5.1	0	0.0 to 0.1
	Both	5 to 9	3.7	2.9 to 4.8	0	0.0 to 0.0
	Both	10 to 14	7.5	5.7 to 9.4	0.1	0.0 to 0.1
	Both	15 to 19	16.8	12.1 to 21.1	0.2	0.1 to 0.2
Western Europe	Male	1 to 4	8.2	5.2 to 11.3	0.1	0.1 to 0.1
	Male	5 to 9	7.1	5.3 to 9.1	0.1	0.0 to 0.1
	Male	10 to 14	13.8	10.6 to 17.0	0.2	0.1 to 0.2
	Male	15 to 19	24.9	19.1 to 31.2	0.3	0.2 to 0.4
	Female	1 to 4	8.1	6.0 to 12.1	0.1	0.1 to 0.1
	Female	5 to 9	7.4	5.7 to 11.3	0.1	0.1 to 0.1
	Female	10 to 14	13.8	10.7 to 17.8	0.2	0.1 to 0.2
	Female	15 to 19	21.7	16.3 to 28.2	0.3	0.2 to 0.4
	Both	1 to 4	8.1	6.1 to 10.7	0.1	0.1 to 0.1

	Both	5 to 9	7.2	5.8 to 9.2	0.1	0.1 to 0.1
	Both	10 to 14	13.8	11.3 to 16.2	0.2	0.1 to 0.2
	Both	15 to 19	23.3	19.1 to 27.6	0.3	0.2 to 0.3
Southern Latin America	Male	1 to 4	33.3	19.2 to 54.7	0.4	0.2 to 0.6
	Male	5 to 9	28.2	19.5 to 38.6	0.3	0.2 to 0.5
	Male	10 to 14	41.7	30.4 to 57.1	0.5	0.4 to 0.7
	Male	15 to 19	100.6	72.8 to 138.3	1.4	1.0 to 2.0
	Female	1 to 4	21.2	12.4 to 37.0	0.2	0.1 to 0.4
	Female	5 to 9	21.8	14.7 to 34.0	0.3	0.2 to 0.4
	Female	10 to 14	34.9	25.9 to 51.5	0.5	0.3 to 0.7
	Female	15 to 19	61.9	43.8 to 88.5	0.9	0.6 to 1.3
	Both	1 to 4	27.3	17.0 to 42.2	0.3	0.2 to 0.5
	Both	5 to 9	25.1	17.9 to 33.7	0.3	0.2 to 0.4
	Both	10 to 14	38.3	30.5 to 49.4	0.5	0.4 to 0.6
	Both	15 to 19	81.6	63.7 to 104.5	1.1	0.9 to 1.5
High-income North America	Male	1 to 4	23.2	12.6 to 37.2	0.3	0.1 to 0.4
	Male	5 to 9	15.2	10.7 to 20.5	0.1	0.1 to 0.2
	Male	10 to 14	24.4	17.3 to 32.0	0.2	0.1 to 0.3
	Male	15 to 19	42	29.8 to 57.9	0.4	0.2 to 0.6
	Female	1 to 4	21.4	11.4 to 32.0	0.2	0.1 to 0.4
	Female	5 to 9	13.5	8.7 to 18.2	0.1	0.1 to 0.2
	Female	10 to 14	20.4	14.7 to 26.7	0.2	0.1 to 0.2
	Female	15 to 19	34.5	23.0 to 45.9	0.3	0.1 to 0.4
	Both	1 to 4	22.3	13.5 to 32.6	0.3	0.1 to 0.4
	Both	5 to 9	14.4	10.7 to 18.3	0.1	0.1 to 0.2
	Both	10 to 14	22.5	17.2 to 28.0	0.2	0.1 to 0.2
	Both	15 to 19	38.3	29.1 to 48.7	0.3	0.2 to 0.4
Caribbean	Male	1 to 4	301.8	150.8 to 534.4	3.6	1.8 to 6.3

	Male	5 to 9	110.2	68.3 to 168.0	1.4	0.9 to 2.1
	Male	10 to 14	104.3	71.7 to 144.6	1.4	0.9 to 1.9
	Male	15 to 19	117	81.6 to 163.5	1.7	1.1 to 2.3
	Female	1 to 4	284.3	155.5 to 481.9	3.4	1.8 to 5.7
	Female	5 to 9	101.5	61.6 to 157.8	1.3	0.8 to 2.0
	Female	10 to 14	89.4	57.0 to 129.8	1.2	0.8 to 1.7
	Female	15 to 19	109.7	73.4 to 159.0	1.6	1.0 to 2.3
	Both	1 to 4	293.2	181.2 to 450.3	3.5	2.1 to 5.3
	Both	5 to 9	105.9	74.1 to 147.1	1.3	0.9 to 1.8
	Both	10 to 14	97	73.8 to 128.7	1.3	1.0 to 1.7
	Both	15 to 19	113.4	86.7 to 148.4	1.6	1.2 to 2.1
Andean Latin America	Male	1 to 4	86.8	50.8 to 147.7	1	0.6 to 1.7
	Male	5 to 9	60.4	42.2 to 84.6	0.8	0.5 to 1.1
	Male	10 to 14	98.7	73.2 to 131.7	1.3	1.0 to 1.8
	Male	15 to 19	171.7	119.6 to 248.2	2.5	1.7 to 3.6
	Female	1 to 4	63.9	36.3 to 125.4	0.8	0.4 to 1.5
	Female	5 to 9	44.4	27.1 to 63.2	0.6	0.3 to 0.8
	Female	10 to 14	77.4	52.5 to 104.4	1	0.7 to 1.4
	Female	15 to 19	122.5	77.3 to 164.5	1.8	1.1 to 2.4
	Both	1 to 4	75.6	50.4 to 119.1	0.9	0.6 to 1.4
	Both	5 to 9	52.6	40.2 to 68.1	0.7	0.5 to 0.9
	Both	10 to 14	88.2	71.1 to 108.7	1.2	0.9 to 1.5
	Both	15 to 19	147.5	115.1 to 191.0	2.1	1.6 to 2.7
Central Latin America	Male	1 to 4	56.3	39.5 to 75.9	0.7	0.5 to 0.9
	Male	5 to 9	31.7	24.5 to 39.9	0.4	0.3 to 0.5
	Male	10 to 14	54.9	43.8 to 67.9	0.7	0.6 to 0.9
	Male	15 to 19	100.7	78.0 to 132.7	1.4	1.1 to 1.9
	Female	1 to 4	47.4	27.6 to 66.1	0.6	0.3 to 0.8

	Female	5 to 9	26.4	17.9 to 34.0	0.3	0.2 to 0.4
	Female	10 to 14	45.8	33.8 to 58.7	0.6	0.4 to 0.8
	Female	15 to 19	78.9	57.9 to 108.6	1.1	0.8 to 1.5
	Both	1 to 4	51.9	39.2 to 65.2	0.6	0.5 to 0.8
	Both	5 to 9	29.1	24.0 to 35.1	0.4	0.3 to 0.4
	Both	10 to 14	50.5	42.7 to 60.2	0.7	0.6 to 0.8
	Both	15 to 19	90	75.9 to 109.8	1.3	1.1 to 1.6
Tropical Latin America	Male	1 to 4	32.9	15.3 to 59.2	0.4	0.2 to 0.7
	Male	5 to 9	31.9	22.9 to 43.4	0.4	0.3 to 0.5
	Male	10 to 14	51.5	35.7 to 69.3	0.7	0.5 to 0.9
	Male	15 to 19	99.1	64.9 to 146.2	1.4	0.9 to 2.1
	Female	1 to 4	22.8	11.8 to 37.9	0.3	0.1 to 0.4
	Female	5 to 9	26.6	19.3 to 38.9	0.3	0.2 to 0.5
	Female	10 to 14	46.6	33.7 to 69.9	0.6	0.4 to 0.9
	Female	15 to 19	69.9	48.3 to 110.5	1	0.7 to 1.6
	Both	1 to 4	27.9	16.5 to 44.9	0.3	0.2 to 0.5
	Both	5 to 9	29.3	23.0 to 37.3	0.4	0.3 to 0.5
	Both	10 to 14	49.1	38.2 to 63.6	0.6	0.5 to 0.8
	Both	15 to 19	84.8	63.4 to 112.7	1.2	0.9 to 1.6
North Africa and Middle East	Male	1 to 4	250.8	176.0 to 344.1	3	2.1 to 4.1
	Male	5 to 9	169.9	137.9 to 207.2	2.1	1.7 to 2.6
	Male	10 to 14	187.2	155.2 to 224.0	2.5	2.1 to 3.0
	Male	15 to 19	311.7	248.9 to 383.2	4.5	3.6 to 5.5
	Female	1 to 4	223.5	161.5 to 310.9	2.6	1.9 to 3.7
	Female	5 to 9	124.7	97.7 to 171.1	1.6	1.2 to 2.2
	Female	10 to 14	123.4	97.3 to 164.9	1.7	1.3 to 2.2
	Female	15 to 19	155.7	117.9 to 202.9	2.2	1.7 to 2.9
	Both	1 to 4	237.5	184.7 to 304.8	2.8	2.2 to 3.6

	Both	5 to 9	147.8	123.2 to 175.5	1.9	1.5 to 2.2
	Both	10 to 14	156.1	133.5 to 184.2	2.1	1.8 to 2.5
	Both	15 to 19	235.7	199.1 to 276.4	3.4	2.9 to 4.0
South Asia	Male	1 to 4	75.4	40.2 to 132.5	0.9	0.5 to 1.6
	Male	5 to 9	54.6	32.3 to 89.4	0.7	0.4 to 1.1
	Male	10 to 14	107.8	71.8 to 159.6	1.4	0.9 to 2.1
	Male	15 to 19	172.7	114.0 to 254.2	2.4	1.6 to 3.6
	Female	1 to 4	56.7	25.9 to 131.4	0.7	0.3 to 1.6
	Female	5 to 9	81.6	45.5 to 155.3	1	0.6 to 1.9
	Female	10 to 14	85.6	50.7 to 147.1	1.1	0.7 to 2.0
	Female	15 to 19	129	78.8 to 203.1	1.8	1.1 to 2.9
	Both	1 to 4	66.4	39.2 to 112.7	0.8	0.5 to 1.3
	Both	5 to 9	67.4	44.4 to 107.1	0.8	0.5 to 1.3
	Both	10 to 14	97.2	68.7 to 136.4	1.3	0.9 to 1.8
	Both	15 to 19	151.8	110.9 to 205.8	2.2	1.6 to 2.9
Central Sub-Saharan Africa	Male	1 to 4	396.9	191.5 to 766.9	4.7	2.3 to 9.1
	Male	5 to 9	96.6	48.2 to 173.9	1.2	0.6 to 2.2
	Male	10 to 14	93.3	52.9 to 148.8	1.2	0.7 to 2.0
	Male	15 to 19	165.5	100.1 to 238.2	2.4	1.4 to 3.4
	Female	1 to 4	316.5	92.9 to 758.8	3.7	1.1 to 9.0
	Female	5 to 9	121.9	60.3 to 221.5	1.5	0.8 to 2.8
	Female	10 to 14	104.1	58.8 to 169.8	1.4	0.8 to 2.3
	Female	15 to 19	154.9	95.1 to 245.0	2.2	1.4 to 3.5
	Both	1 to 4	357.1	178.2 to 653.9	4.2	2.1 to 7.7
	Both	5 to 9	109.2	65.1 to 178.3	1.4	0.8 to 2.2
	Both	10 to 14	98.7	65.4 to 141.3	1.3	0.9 to 1.9
	Both	15 to 19	160.2	112.3 to 217.8	2.3	1.6 to 3.1
Eastern Sub-Saharan Africa	Male	1 to 4	20.9	12.7 to 33.8	0.2	0.1 to 0.4

	Male	5 to 9	18	11.1 to 26.1	0.2	0.1 to 0.3
	Male	10 to 14	24.7	16.5 to 34.6	0.3	0.2 to 0.4
	Male	15 to 19	96.6	68.8 to 128.6	1.4	1.0 to 1.8
	Female	1 to 4	15.7	8.4 to 32.6	0.2	0.1 to 0.4
	Female	5 to 9	11.7	7.3 to 20.1	0.1	0.1 to 0.2
	Female	10 to 14	13.7	8.8 to 22.1	0.2	0.1 to 0.3
	Female	15 to 19	44.6	26.2 to 65.3	0.6	0.4 to 0.9
	Both	1 to 4	18.3	12.0 to 27.7	0.2	0.1 to 0.3
	Both	5 to 9	14.9	10.4 to 20.0	0.2	0.1 to 0.2
	Both	10 to 14	19.3	13.7 to 25.6	0.2	0.2 to 0.3
	Both	15 to 19	70.7	51.8 to 90.0	1	0.7 to 1.3
Southern Sub-Saharan Africa	Male	1 to 4	165.5	72.8 to 321.0	2	0.9 to 3.8
	Male	5 to 9	46.9	25.6 to 78.9	0.6	0.3 to 1.0
	Male	10 to 14	43.5	28.0 to 65.9	0.6	0.4 to 0.9
	Male	15 to 19	121.3	80.7 to 169.6	1.7	1.1 to 2.4
	Female	1 to 4	301.1	63.1 to 646.9	3.6	0.7 to 7.7
	Female	5 to 9	59.7	23.2 to 118.0	0.7	0.3 to 1.5
	Female	10 to 14	38.8	19.7 to 63.7	0.5	0.3 to 0.8
	Female	15 to 19	79.3	46.0 to 116.8	1.1	0.6 to 1.7
	Both	1 to 4	232.6	82.9 to 423.2	2.8	1.0 to 5.0
	Both	5 to 9	53.3	27.4 to 87.8	0.7	0.3 to 1.1
	Both	10 to 14	41.1	26.4 to 58.3	0.5	0.3 to 0.8
	Both	15 to 19	100.4	71.1 to 132.7	1.4	1.0 to 1.9
Western Sub-Saharan Africa	Male	1 to 4	201.9	106.1 to 373.7	2.4	1.3 to 4.4
	Male	5 to 9	77.7	46.9 to 127.0	1	0.6 to 1.6
	Male	10 to 14	70.2	44.8 to 104.9	0.9	0.6 to 1.4
	Male	15 to 19	92.2	65.1 to 123.3	1.3	0.9 to 1.8
	Female	1 to 4	182.2	79.5 to 522.7	2.2	0.9 to 6.2

	Female	5 to 9	51.5	31.2 to 98.7	0.6	0.4 to 1.2
	Female	10 to 14	50.1	32.3 to 94.7	0.7	0.4 to 1.3
	Female	15 to 19	81.4	54.8 to 120.8	1.2	0.8 to 1.7
	Both	1 to 4	192.2	113.0 to 367.9	2.3	1.3 to 4.4
	Both	5 to 9	64.8	43.0 to 97.8	0.8	0.5 to 1.2
	Both	10 to 14	60.3	43.2 to 84.3	0.8	0.6 to 1.1
	Both	15 to 19	86.9	66.5 to 111.6	1.2	0.9 to 1.6

Ischemic Stroke								
East Asia	Male	1 to 4	0.3	0.3 to 0.4	8.2	4.2 to 14.1	0.1	0.0 to 0.2
	Male	5 to 9	0.9	0.8 to 1.1	5.5	3.2 to 8.3	0.1	0.0 to 0.1
	Male	10 to 14	2	1.8 to 2.2	6.8	4.4 to 10.0	0.1	0.1 to 0.1
	Male	15 to 19	3.3	2.9 to 3.8	16.2	10.7 to 24.3	0.2	0.1 to 0.3
	Female	1 to 4	0.2	0.2 to 0.2	7.5	3.4 to 12.4	0.1	0.0 to 0.1
	Female	5 to 9	0.7	0.6 to 0.8	6.8	3.8 to 9.4	0.1	0.0 to 0.1
	Female	10 to 14	1.5	1.3 to 1.7	4.9	2.7 to 6.7	0.1	0.0 to 0.1
	Female	15 to 19	2.6	2.3 to 3.0	7	3.7 to 10.5	0.1	0.0 to 0.1
	Both	1 to 4	0.3	0.2 to 0.3	7.9	4.4 to 12.7	0.1	0.1 to 0.1
	Both	5 to 9	0.8	0.7 to 0.9	6.1	4.2 to 8.3	0.1	0.1 to 0.1
	Both	10 to 14	1.8	1.6 to 1.9	5.9	4.3 to 8.0	0.1	0.1 to 0.1
	Both	15 to 19	3	2.7 to 3.3	11.9	8.3 to 16.2	0.2	0.1 to 0.2
Southeast Asia	Male	1 to 4	0.4	0.3 to 0.5	8.5	4.7 to 13.4	0.1	0.1 to 0.2
	Male	5 to 9	1.1	0.8 to 1.5	3.9	2.2 to 5.8	0	0.0 to 0.1
	Male	10 to 14	2.3	1.8 to 2.9	3.4	2.3 to 5.1	0	0.0 to 0.1
	Male	15 to 19	4	3.2 to 5.0	21.9	12.8 to 33.2	0.3	0.2 to 0.5
	Female	1 to 4	0.2	0.2 to 0.3	5.2	2.5 to 8.1	0.1	0.0 to 0.1
	Female	5 to 9	0.8	0.6 to 1.0	4.1	2.4 to 6.0	0	0.0 to 0.1
	Female	10 to 14	1.7	1.3 to 2.1	3.9	2.5 to 7.0	0	0.0 to 0.1
	Female	15 to 19	3	2.3 to 3.8	12.5	7.7 to 18.7	0.2	0.1 to 0.3
	Both	1 to 4	0.3	0.2 to 0.4	6.9	4.1 to 10.0	0.1	0.0 to 0.1
	Both	5 to 9	0.9	0.7 to 1.2	4	2.7 to 5.3	0	0.0 to 0.1
	Both	10 to 14	2	1.7 to 2.4	3.6	2.7 to 5.4	0	0.0 to 0.1
	Both	15 to 19	3.5	3.0 to 4.1	17.3	11.8 to 23.4	0.2	0.2 to 0.3
Oceania	Male	1 to 4	0.3	0.2 to 0.5	22.7	8.6 to 54.6	0.3	0.1 to 0.6

	Male	5 to 9	1	0.6 to 1.5	24.7	11.7 to 44.1	0.3	0.1 to 0.6
	Male	10 to 14	2.1	1.4 to 2.9	7.9	4.1 to 12.8	0.1	0.1 to 0.2
	Male	15 to 19	3.5	2.4 to 4.9	27.3	14.9 to 48.0	0.4	0.2 to 0.7
	Female	1 to 4	0.2	0.1 to 0.3	8.6	2.7 to 26.7	0.1	0.0 to 0.3
	Female	5 to 9	0.7	0.4 to 1.0	3.3	1.3 to 8.1	0	0.0 to 0.1
	Female	10 to 14	1.4	1.0 to 2.1	10.3	4.7 to 20.9	0.1	0.1 to 0.3
	Female	15 to 19	2.5	1.7 to 3.6	5	2.7 to 10.0	0.1	0.0 to 0.1
	Both	1 to 4	0.3	0.2 to 0.4	15.9	6.6 to 36.8	0.2	0.1 to 0.4
	Both	5 to 9	0.8	0.6 to 1.2	14.4	7.4 to 25.0	0.2	0.1 to 0.3
	Both	10 to 14	1.8	1.3 to 2.4	9.1	5.3 to 15.2	0.1	0.1 to 0.2
	Both	15 to 19	3	2.3 to 3.9	16.5	9.8 to 27.6	0.2	0.1 to 0.4
Central Asia	Male	1 to 4	0.3	0.3 to 0.4	4.5	2.7 to 7.6	0.1	0.0 to 0.1
	Male	5 to 9	1	0.7 to 1.2	7.1	4.8 to 10.2	0.1	0.1 to 0.1
	Male	10 to 14	1.8	1.4 to 2.2	6.7	4.6 to 9.6	0.1	0.1 to 0.1
	Male	15 to 19	2.7	2.1 to 3.5	25.1	15.5 to 38.2	0.4	0.2 to 0.5
	Female	1 to 4	0.3	0.2 to 0.3	7.6	4.1 to 12.2	0.1	0.0 to 0.1
	Female	5 to 9	0.8	0.6 to 1.0	7.1	4.3 to 9.8	0.1	0.1 to 0.1
	Female	10 to 14	1.6	1.3 to 1.9	4.7	3.2 to 8.2	0.1	0.0 to 0.1
	Female	15 to 19	2.5	2.0 to 3.1	18.4	12.8 to 31.2	0.3	0.2 to 0.4
	Both	1 to 4	0.3	0.2 to 0.4	6	4.2 to 9.2	0.1	0.0 to 0.1
	Both	5 to 9	0.9	0.7 to 1.1	7.1	5.2 to 9.0	0.1	0.1 to 0.1
	Both	10 to 14	1.7	1.4 to 2.0	5.8	4.4 to 7.9	0.1	0.1 to 0.1
	Both	15 to 19	2.6	2.2 to 3.1	21.8	16.1 to 29.9	0.3	0.2 to 0.4
Central Europe	Male	1 to 4	1.3	1.1 to 1.5	3.9	2.2 to 6.1	0	0.0 to 0.1
	Male	5 to 9	3.8	3.3 to 4.5	3.1	2.0 to 4.3	0	0.0 to 0.0
	Male	10 to 14	7.3	6.3 to 8.5	4	2.8 to 5.4	0	0.0 to 0.1
	Male	15 to 19	11.2	9.5 to 13.1	10.2	6.6 to 13.7	0.1	0.1 to 0.2

	Female	1 to 4	0.8	0.6 to 0.9	4.6	2.2 to 7.3	0.1	0.0 to 0.1
	Female	5 to 9	2.3	2.0 to 2.7	3.2	1.8 to 4.4	0	0.0 to 0.1
	Female	10 to 14	4.5	3.9 to 5.2	4.6	3.0 to 6.1	0.1	0.0 to 0.1
	Female	15 to 19	6.9	5.7 to 8.2	8.3	5.0 to 10.6	0.1	0.1 to 0.1
	Both	1 to 4	1	0.9 to 1.2	4.2	2.4 to 6.2	0	0.0 to 0.1
	Both	5 to 9	3.1	2.8 to 3.5	3.1	2.1 to 4.2	0	0.0 to 0.0
	Both	10 to 14	6	5.3 to 6.7	4.3	3.3 to 5.4	0	0.0 to 0.1
	Both	15 to 19	9.1	8.0 to 10.3	9.3	6.5 to 11.6	0.1	0.1 to 0.1
Eastern Europe	Male	1 to 4	3	2.3 to 3.8	1.3	0.8 to 2.0	0	0.0 to 0.0
	Male	5 to 9	8.2	6.4 to 10.0	1.8	1.3 to 2.5	0	0.0 to 0.0
	Male	10 to 14	15	12.0 to 18.3	3	2.0 to 4.1	0	0.0 to 0.0
	Male	15 to 19	22.4	17.8 to 27.9	10.3	6.6 to 15.6	0.1	0.1 to 0.2
	Female	1 to 4	1.8	1.3 to 2.2	1.1	0.6 to 1.7	0	0.0 to 0.0
	Female	5 to 9	5	3.9 to 6.1	1.5	1.0 to 2.0	0	0.0 to 0.0
	Female	10 to 14	9.6	7.5 to 11.7	2.5	1.7 to 3.4	0	0.0 to 0.0
	Female	15 to 19	14.8	11.6 to 18.6	8.7	5.3 to 12.3	0.1	0.0 to 0.1
	Both	1 to 4	2.4	2.0 to 2.8	1.2	0.8 to 1.7	0	0.0 to 0.0
	Both	5 to 9	6.6	5.5 to 7.7	1.7	1.2 to 2.2	0	0.0 to 0.0
	Both	10 to 14	12.4	10.5 to 14.3	2.7	2.0 to 3.6	0	0.0 to 0.0
	Both	15 to 19	18.7	15.7 to 22.1	9.5	7.0 to 13.0	0.1	0.1 to 0.1
High-income Asia Pacific	Male	1 to 4	1.1	0.7 to 1.7	3.1	1.4 to 5.2	0	0.0 to 0.1
	Male	5 to 9	3.4	2.3 to 4.9	1.2	0.7 to 1.6	0	0.0 to 0.0
	Male	10 to 14	6.9	5.1 to 9.1	2.2	1.5 to 3.1	0	0.0 to 0.0

	Male	15 to 19	11.2	8.3 to 15.1	4.8	3.1 to 6.6	0	0.0 to 0.1
	Female	1 to 4	0.7	0.4 to 1.0	1.4	0.8 to 2.6	0	0.0 to 0.0
	Female	5 to 9	2.2	1.5 to 3.1	1.6	0.8 to 2.2	0	0.0 to 0.0
	Female	10 to 14	4.7	3.4 to 6.3	1.9	1.3 to 2.8	0	0.0 to 0.0
	Female	15 to 19	8	6.0 to 10.6	3.5	2.4 to 5.9	0	0.0 to 0.1
	Both	1 to 4	0.9	0.6 to 1.2	2.3	1.3 to 3.4	0	0.0 to 0.0
	Both	5 to 9	2.8	2.1 to 3.7	1.4	0.9 to 1.8	0	0.0 to 0.0
	Both	10 to 14	5.8	4.6 to 7.4	2.1	1.5 to 2.8	0	0.0 to 0.0
	Both	15 to 19	9.7	7.8 to 12.0	4.2	3.0 to 5.6	0	0.0 to 0.1
Australasia	Male	1 to 4	1.2	0.8 to 1.8	1.5	0.8 to 2.4	0	0.0 to 0.0
	Male	5 to 9	3.9	2.6 to 5.4	1	0.7 to 1.4	0	0.0 to 0.0
	Male	10 to 14	8	5.6 to 10.9	1.8	1.2 to 2.7	0	0.0 to 0.0
	Male	15 to 19	13	8.7 to 18.1	3.4	2.2 to 4.8	0	0.0 to 0.0
	Female	1 to 4	0.9	0.6 to 1.3	0.7	0.4 to 1.1	0	0.0 to 0.0
	Female	5 to 9	2.9	2.0 to 4.1	1	0.6 to 1.4	0	0.0 to 0.0
	Female	10 to 14	6.2	4.5 to 8.2	1.7	1.1 to 2.3	0	0.0 to 0.0
	Female	15 to 19	10.4	7.5 to 13.8	3.2	2.2 to 4.4	0	0.0 to 0.0
	Both	1 to 4	1.1	0.8 to 1.4	1.1	0.7 to 1.6	0	0.0 to 0.0
	Both	5 to 9	3.4	2.6 to 4.4	1	0.7 to 1.3	0	0.0 to 0.0
	Both	10 to 14	7.1	5.6 to 8.8	1.8	1.3 to 2.3	0	0.0 to 0.0
	Both	15 to 19	11.7	9.0 to 14.9	3.3	2.4 to 4.4	0	0.0 to 0.0
Western Europe	Male	1 to 4	1.6	1.4 to 1.9	1.3	0.8 to 1.8	0	0.0 to 0.0
	Male	5 to 9	4.7	4.1 to 5.3	1.4	1.0 to 1.9	0	0.0 to 0.0

	Male	10 to 14	9.2	8.1 to 10.3	2.5	1.9 to 3.2	0	0.0 to 0.0
	Male	15 to 19	14.7	13.1 to 16.5	4.5	3.3 to 5.8	0	0.0 to 0.0
	Female	1 to 4	1.3	1.2 to 1.5	1.5	1.1 to 2.7	0	0.0 to 0.0
	Female	5 to 9	4.2	3.7 to 4.7	1.5	1.1 to 2.1	0	0.0 to 0.0
	Female	10 to 14	8.7	7.8 to 9.6	2.5	1.9 to 3.4	0	0.0 to 0.0
	Female	15 to 19	14.3	12.9 to 15.9	4.6	3.4 to 5.9	0	0.0 to 0.0
	Both	1 to 4	1.5	1.3 to 1.6	1.4	1.0 to 2.0	0	0.0 to 0.0
	Both	5 to 9	4.4	4.0 to 4.8	1.4	1.1 to 1.9	0	0.0 to 0.0
	Both	10 to 14	8.9	8.3 to 9.6	2.5	1.9 to 3.2	0	0.0 to 0.0
	Both	15 to 19	14.5	13.4 to 15.7	4.5	3.5 to 5.7	0	0.0 to 0.0
Southern Latin America	Male	1 to 4	0.7	0.5 to 1.1	7.4	3.8 to 14.3	0.1	0.0 to 0.2
	Male	5 to 9	2.2	1.5 to 3.0	5.8	3.5 to 10.0	0.1	0.0 to 0.1
	Male	10 to 14	4.4	3.3 to 5.8	7.4	4.6 to 12.2	0.1	0.1 to 0.2
	Male	15 to 19	7.2	5.2 to 9.5	21.5	13.4 to 36.0	0.3	0.2 to 0.5
	Female	1 to 4	0.6	0.4 to 0.9	5.9	3.2 to 12.0	0.1	0.0 to 0.1
	Female	5 to 9	1.8	1.2 to 2.6	5.1	3.0 to 10.2	0.1	0.0 to 0.1
	Female	10 to 14	3.7	2.6 to 4.9	6.5	4.1 to 12.6	0.1	0.0 to 0.2
	Female	15 to 19	5.9	4.5 to 7.8	12.1	8.1 to 20.9	0.2	0.1 to 0.3
	Both	1 to 4	0.7	0.5 to 0.9	6.7	3.8 to 12.1	0.1	0.0 to 0.1
	Both	5 to 9	2	1.6 to 2.6	5.5	3.6 to 9.2	0.1	0.0 to 0.1
	Both	10 to 14	4	3.3 to 5.0	6.9	4.8 to 11.3	0.1	0.1 to 0.1
	Both	15 to 19	6.6	5.3 to 8.0	16.9	11.8 to 26.4	0.2	0.2 to 0.4
High-income North America	Male	1 to 4	4.5	3.2 to 5.9	5.1	2.5 to 9.1	0.1	0.0 to 0.1
	Male	5 to 9	16.8	12.6 to 20.9	4.1	2.7 to 5.7	0	0.0 to 0.0

	Male	10 to 14	38.5	30.1 to 47.6	7.5	4.4 to 11.2	0	0.0 to 0.0
	Male	15 to 19	67.9	52.3 to 84.1	13.3	7.3 to 21.6	0	0.0 to 0.1
	Female	1 to 4	3.2	1.9 to 4.3	4.7	2.6 to 7.9	0	0.0 to 0.1
	Female	5 to 9	12.8	8.4 to 16.6	3.5	2.3 to 4.9	0	0.0 to 0.0
	Female	10 to 14	31.4	22.8 to 39.2	6.3	3.6 to 9.8	0	0.0 to 0.0
	Female	15 to 19	58.1	42.9 to 73.0	11.9	6.6 to 19.1	0	0.0 to 0.1
	Both	1 to 4	3.8	3.0 to 4.7	4.9	3.0 to 7.9	0.1	0.0 to 0.1
	Both	5 to 9	14.8	11.9 to 17.7	3.8	2.7 to 5.1	0	0.0 to 0.0
	Both	10 to 14	35	29.9 to 41.0	6.9	4.5 to 10.0	0	0.0 to 0.0
	Both	15 to 19	63.1	53.7 to 75.1	12.6	8.0 to 19.0	0	0.0 to 0.1
Caribbean	Male	1 to 4	0.5	0.4 to 0.7	49.3	23.2 to 96.5	0.6	0.3 to 1.1
	Male	5 to 9	1.7	1.3 to 2.2	20.8	11.7 to 36.1	0.3	0.1 to 0.5
	Male	10 to 14	3.8	2.9 to 4.6	23.7	14.2 to 36.7	0.3	0.2 to 0.5
	Male	15 to 19	6.6	5.0 to 8.1	35.9	22.1 to 55.4	0.5	0.3 to 0.8
	Female	1 to 4	0.3	0.2 to 0.4	46.5	22.0 to 88.8	0.5	0.3 to 1.1
	Female	5 to 9	0.9	0.6 to 1.3	18.8	9.1 to 32.4	0.2	0.1 to 0.4
	Female	10 to 14	2.3	1.8 to 2.9	20.1	11.1 to 36.8	0.3	0.1 to 0.5
	Female	15 to 19	4.4	3.5 to 5.4	30.3	17.9 to 51.7	0.4	0.2 to 0.7
	Both	1 to 4	0.4	0.3 to 0.5	47.9	26.4 to 80.7	0.6	0.3 to 1.0
	Both	5 to 9	1.3	1.0 to 1.6	19.8	12.8 to 31.4	0.2	0.2 to 0.4
	Both	10 to 14	3.1	2.6 to 3.6	21.9	15.0 to 33.0	0.3	0.2 to 0.4
	Both	15 to 19	5.5	4.7 to 6.5	33.1	23.3 to 48.4	0.5	0.3 to 0.7
Andean Latin America	Male	1 to 4	0.4	0.3 to 0.6	12.7	6.3 to 24.4	0.1	0.1 to 0.3

	Male	5 to 9	1.4	0.9 to 1.9	13.2	7.6 to 22.9	0.2	0.1 to 0.3
	Male	10 to 14	2.9	2.1 to 3.7	17.9	10.4 to 30.6	0.2	0.1 to 0.4
	Male	15 to 19	4.6	3.3 to 6.0	42.1	24.5 to 73.4	0.6	0.3 to 1.0
	Female	1 to 4	0.2	0.1 to 0.4	8.9	4.4 to 19.8	0.1	0.1 to 0.2
	Female	5 to 9	0.8	0.4 to 1.1	10.5	6.2 to 20.1	0.1	0.1 to 0.3
	Female	10 to 14	1.7	1.2 to 2.2	16.7	10.6 to 28.6	0.2	0.1 to 0.4
	Female	15 to 19	2.7	2.0 to 3.6	29.9	18.7 to 50.7	0.4	0.3 to 0.7
	Both	1 to 4	0.3	0.3 to 0.5	10.8	6.3 to 21.2	0.1	0.1 to 0.3
	Both	5 to 9	1.1	0.8 to 1.4	11.9	7.9 to 19.8	0.1	0.1 to 0.2
	Both	10 to 14	2.3	1.9 to 2.8	17.3	12.2 to 26.3	0.2	0.2 to 0.3
	Both	15 to 19	3.6	3.0 to 4.5	36.1	24.8 to 58.3	0.5	0.4 to 0.8
Central Latin America	Male	1 to 4	0.1	0.0 to 0.1	9.7	5.1 to 17.2	0.1	0.1 to 0.2
	Male	5 to 9	0.2	0.2 to 0.3	5.5	3.3 to 8.9	0.1	0.0 to 0.1
	Male	10 to 14	0.5	0.4 to 0.6	7.5	4.9 to 12.2	0.1	0.1 to 0.2
	Male	15 to 19	1	0.8 to 1.2	17.4	11.7 to 28.1	0.2	0.2 to 0.4
	Female	1 to 4	0.1	0.0 to 0.1	7.7	3.5 to 15.8	0.1	0.0 to 0.2
	Female	5 to 9	0.2	0.2 to 0.2	4.6	2.6 to 7.8	0.1	0.0 to 0.1
	Female	10 to 14	0.4	0.4 to 0.5	6.8	4.5 to 11.4	0.1	0.1 to 0.2
	Female	15 to 19	0.8	0.7 to 0.9	13.8	9.5 to 21.8	0.2	0.1 to 0.3
	Both	1 to 4	0.1	0.1 to 0.1	8.7	5.3 to 15.1	0.1	0.1 to 0.2
	Both	5 to 9	0.2	0.2 to 0.2	5	3.5 to 7.7	0.1	0.0 to 0.1
	Both	10 to 14	0.5	0.4 to 0.5	7.2	5.4 to 10.9	0.1	0.1 to 0.1
	Both	15 to 19	0.9	0.8 to 1.0	15.6	11.9 to 23.5	0.2	0.2 to 0.3
Tropical Latin America	Male	1 to 4	0.8	0.5 to 1.2	4.3	1.5 to 9.0	0	0.0 to 0.1
	Male	5 to 9	2.5	1.5 to 3.7	3.3	1.9 to 5.1	0	0.0 to 0.1
	Male	10 to 14	5.3	3.4 to 7.3	4.2	2.7 to 6.4	0	0.0 to 0.1
	Male	15 to 19	8.8	5.5 to 12.3	14.2	8.2 to 22.7	0.2	0.1 to 0.3

	Female	1 to 4	0.5	0.3 to 0.8	2.6	1.1 to 5.0	0	0.0 to 0.1
	Female	5 to 9	1.6	0.8 to 2.4	2.7	1.5 to 3.9	0	0.0 to 0.0
	Female	10 to 14	3.6	2.3 to 5.1	4.1	2.6 to 6.2	0	0.0 to 0.1
	Female	15 to 19	6.2	3.8 to 8.9	9.9	6.0 to 15.9	0.1	0.1 to 0.2
	Both	1 to 4	0.6	0.4 to 0.9	3.5	1.5 to 6.7	0	0.0 to 0.1
	Both	5 to 9	2.1	1.4 to 2.7	3	2.0 to 4.1	0	0.0 to 0.0
	Both	10 to 14	4.4	3.3 to 5.6	4.2	3.0 to 5.6	0	0.0 to 0.1
	Both	15 to 19	7.5	5.5 to 9.9	12.1	8.3 to 16.8	0.2	0.1 to 0.2
North Africa and Middle East	Male	1 to 4	0.5	0.4 to 0.6	23	13.9 to 37.4	0.3	0.2 to 0.4
	Male	5 to 9	1.6	1.4 to 1.8	11.2	7.8 to 16.3	0.1	0.1 to 0.2
	Male	10 to 14	3.1	2.7 to 3.4	15.8	9.7 to 22.0	0.2	0.1 to 0.3
	Male	15 to 19	4.7	4.1 to 5.3	32.5	23.3 to 47.6	0.5	0.3 to 0.7
	Female	1 to 4	0.3	0.3 to 0.4	28.3	16.3 to 45.8	0.3	0.2 to 0.5
	Female	5 to 9	1	0.9 to 1.1	10.9	7.2 to 16.1	0.1	0.1 to 0.2
	Female	10 to 14	2	1.8 to 2.2	15.6	9.5 to 23.1	0.2	0.1 to 0.3
	Female	15 to 19	3.2	2.8 to 3.7	20.4	13.5 to 29.5	0.3	0.2 to 0.4
	Both	1 to 4	0.4	0.4 to 0.5	25.6	17.4 to 36.3	0.3	0.2 to 0.4
	Both	5 to 9	1.3	1.2 to 1.4	11.1	8.4 to 14.9	0.1	0.1 to 0.2
	Both	10 to 14	2.5	2.3 to 2.8	15.7	10.9 to 20.3	0.2	0.1 to 0.3
	Both	15 to 19	3.9	3.6 to 4.3	26.6	20.8 to 35.1	0.4	0.3 to 0.5
South Asia	Male	1 to 4	0.8	0.6 to 1.1	14.1	6.0 to 28.5	0.2	0.1 to 0.3
	Male	5 to 9	2.7	1.8 to 3.8	8.4	4.0 to 16.1	0.1	0.0 to 0.2
	Male	10 to 14	6	4.3 to 8.3	13.4	7.1 to 23.3	0.2	0.1 to 0.3
	Male	15 to 19	10.3	7.3 to 14.1	26.6	12.8 to 46.9	0.4	0.2 to 0.7
	Female	1 to 4	0.6	0.4 to 0.8	9.8	3.8 to 21.5	0.1	0.0 to 0.3
	Female	5 to 9	2	1.4 to 2.6	12	5.1 to 25.6	0.1	0.1 to 0.3
	Female	10 to 14	4.3	3.1 to 5.5	10.5	4.7 to 23.1	0.1	0.1 to 0.3

	Female	15 to 19	7.2	5.3 to 9.3	20.3	9.6 to 39.9	0.3	0.1 to 0.6
	Both	1 to 4	0.7	0.5 to 0.9	12.1	6.4 to 21.3	0.1	0.1 to 0.3
	Both	5 to 9	2.4	1.8 to 3.0	10.1	5.5 to 16.9	0.1	0.1 to 0.2
	Both	10 to 14	5.2	4.2 to 6.4	12	7.3 to 18.8	0.2	0.1 to 0.2
	Both	15 to 19	8.8	7.0 to 11.0	23.6	14.3 to 36.1	0.3	0.2 to 0.5
Central Sub-Saharan Africa	Male	1 to 4	0.7	0.5 to 1.0	88.8	10.5 to 274.4	1.1	0.1 to 3.2
	Male	5 to 9	2.1	1.5 to 2.8	15	5.4 to 34.4	0.2	0.1 to 0.4
	Male	10 to 14	4.2	3.1 to 5.4	11.5	5.3 to 22.5	0.1	0.1 to 0.3
	Male	15 to 19	6.5	4.8 to 8.6	24.9	9.9 to 45.8	0.3	0.1 to 0.6
	Female	1 to 4	0.5	0.3 to 0.7	58.4	4.6 to 153.7	0.7	0.1 to 1.8
	Female	5 to 9	1.6	1.1 to 2.0	17.9	4.5 to 39.9	0.2	0.1 to 0.5
	Female	10 to 14	3.1	2.2 to 4.0	11.8	4.8 to 24.7	0.2	0.1 to 0.3
	Female	15 to 19	4.8	3.5 to 6.3	22.4	10.6 to 44.1	0.3	0.1 to 0.6
	Both	1 to 4	0.6	0.5 to 0.8	73.7	8.9 to 173.6	0.9	0.1 to 2.1
	Both	5 to 9	1.8	1.5 to 2.2	16.4	5.6 to 31.7	0.2	0.1 to 0.4
	Both	10 to 14	3.6	2.9 to 4.4	11.6	6.0 to 19.6	0.1	0.1 to 0.3
	Both	15 to 19	5.7	4.5 to 7.1	23.7	13.8 to 37.3	0.3	0.2 to 0.5
Eastern Sub-Saharan Africa	Male	1 to 4	0.8	0.7 to 1.0	3.8	1.1 to 8.0	0	0.0 to 0.1
	Male	5 to 9	2.3	2.0 to 2.6	2.6	1.2 to 4.3	0	0.0 to 0.0
	Male	10 to 14	4	3.5 to 4.6	3.1	1.6 to 5.0	0	0.0 to 0.1
	Male	15 to 19	5.6	4.7 to 6.6	13.7	5.1 to 25.0	0.2	0.1 to 0.3
	Female	1 to 4	0.5	0.5 to 0.6	2.4	1.1 to 5.0	0	0.0 to 0.1
	Female	5 to 9	1.5	1.3 to 1.7	1.7	0.8 to 2.9	0	0.0 to 0.0
	Female	10 to 14	2.7	2.3 to 3.1	1.8	0.9 to 3.0	0	0.0 to 0.0
	Female	15 to 19	3.8	3.2 to 4.4	6.4	2.9 to 10.7	0.1	0.0 to 0.1
	Both	1 to 4	0.7	0.6 to 0.8	3.1	1.4 to 5.3	0	0.0 to 0.1
	Both	5 to 9	1.9	1.7 to 2.1	2.2	1.1 to 3.2	0	0.0 to 0.0

	Both	10 to 14	3.4	3.0 to 3.7	2.4	1.4 to 3.6	0	0.0 to 0.0
	Both	15 to 19	4.7	4.1 to 5.3	10	4.6 to 16.1	0.1	0.1 to 0.2
Southern Sub-Saharan Africa	Male	1 to 4	0.5	0.4 to 0.7	3.5	1.1 to 9.4	0	0.0 to 0.1
	Male	5 to 9	1.7	1.2 to 2.3	1.9	0.8 to 4.1	0	0.0 to 0.0
	Male	10 to 14	3.6	2.7 to 4.7	3.1	1.7 to 5.9	0	0.0 to 0.1
	Male	15 to 19	5.9	4.3 to 7.6	17.1	10.3 to 26.7	0.2	0.1 to 0.4
	Female	1 to 4	0.5	0.4 to 0.7	7.5	2.3 to 23.8	0.1	0.0 to 0.3
	Female	5 to 9	1.6	1.2 to 2.1	2.6	1.1 to 8.0	0	0.0 to 0.1
	Female	10 to 14	3.4	2.6 to 4.3	2.6	1.3 to 6.5	0	0.0 to 0.1
	Female	15 to 19	5.7	4.1 to 7.4	9.2	5.1 to 17.6	0.1	0.1 to 0.2
	Both	1 to 4	0.5	0.4 to 0.6	5.5	2.1 to 15.2	0.1	0.0 to 0.2
	Both	5 to 9	1.7	1.4 to 2.0	2.2	1.2 to 5.8	0	0.0 to 0.1
	Both	10 to 14	3.5	2.8 to 4.2	2.9	1.8 to 5.3	0	0.0 to 0.1
	Both	15 to 19	5.8	4.6 to 7.1	13.2	8.8 to 19.8	0.2	0.1 to 0.3
Western Sub-Saharan Africa	Male	1 to 4	0.7	0.5 to 0.8	35.8	11.8 to 96.9	0.4	0.1 to 1.1
	Male	5 to 9	2	1.7 to 2.5	9	4.5 to 19.0	0.1	0.1 to 0.2
	Male	10 to 14	4.1	3.4 to 4.9	12.1	6.5 to 20.1	0.2	0.1 to 0.3
	Male	15 to 19	6.4	5.2 to 7.9	25.8	12.7 to 39.1	0.4	0.2 to 0.6
	Female	1 to 4	0.5	0.4 to 0.6	41.7	12.5 to 126.6	0.5	0.1 to 1.5
	Female	5 to 9	1.6	1.3 to 2.0	8.7	4.2 to 21.5	0.1	0.0 to 0.3
	Female	10 to 14	3.2	2.7 to 3.9	4.6	2.1 to 13.9	0.1	0.0 to 0.2
	Female	15 to 19	5	4.0 to 6.1	13.3	7.9 to 25.8	0.2	0.1 to 0.4
	Both	1 to 4	0.6	0.5 to 0.7	38.7	14.2 to 89.8	0.5	0.2 to 1.1
	Both	5 to 9	1.8	1.6 to 2.1	8.9	5.0 to 18.4	0.1	0.1 to 0.2
	Both	10 to 14	3.7	3.2 to 4.2	8.4	5.3 to 13.9	0.1	0.1 to 0.2
	Both	15 to 19	5.7	4.9 to 6.6	19.7	12.2 to 27.8	0.3	0.2 to 0.4

Haemorrhagic Stroke								
East Asia	Male	1 to 4	1.2	1.1 to 1.3	71.6	45.8 to 102.7	0.8	0.5 to 1.2
	Male	5 to 9	3.5	3.2 to 3.8	41.8	32.8 to 52.6	0.5	0.4 to 0.7
	Male	10 to 14	6.8	6.3 to 7.3	48.4	38.3 to 60.5	0.6	0.5 to 0.8
	Male	15 to 19	10.6	9.8 to 11.6	145.9	114.8 to 196.6	2.1	1.6 to 2.8
	Female	1 to 4	0.7	0.6 to 0.8	37.3	22.4 to 50.0	0.4	0.3 to 0.6
	Female	5 to 9	2.1	2.0 to 2.3	35.1	25.3 to 44.6	0.4	0.3 to 0.6
	Female	10 to 14	4.3	3.9 to 4.7	38	27.2 to 48.8	0.5	0.4 to 0.6
	Female	15 to 19	6.8	6.2 to 7.5	58.1	39.6 to 74.6	0.8	0.6 to 1.1
	Both	1 to 4	1	0.9 to 1.0	55.7	37.2 to 76.6	0.7	0.4 to 0.9
	Both	5 to 9	2.9	2.7 to 3.0	38.7	31.8 to 46.1	0.5	0.4 to 0.6
	Both	10 to 14	5.6	5.3 to 6.0	43.6	36.5 to 51.8	0.6	0.5 to 0.7
	Both	15 to 19	8.8	8.3 to 9.5	104.8	86.7 to 133.0	1.5	1.2 to 1.9
Southeast Asia	Male	1 to 4	1.3	1.1 to 1.6	50.1	33.0 to 73.2	0.6	0.4 to 0.9
	Male	5 to 9	4	3.4 to 4.8	24.4	16.6 to 35.9	0.3	0.2 to 0.4
	Male	10 to 14	8	6.9 to 9.2	42.6	30.3 to 59.0	0.6	0.4 to 0.8
	Male	15 to 19	12.8	11.0 to 15.0	140.8	104.9 to 185.1	2	1.5 to 2.6
	Female	1 to 4	0.8	0.6 to 0.9	28.1	17.4 to 40.9	0.3	0.2 to 0.5
	Female	5 to 9	2.4	2.0 to 2.8	26.8	18.8 to 42.6	0.3	0.2 to 0.5
	Female	10 to 14	4.8	4.0 to 5.7	34.7	23.8 to 56.9	0.5	0.3 to 0.8
	Female	15 to 19	7.9	6.5 to 9.5	64	45.5 to 90.3	0.9	0.6 to 1.3
	Both	1 to 4	1.1	0.9 to 1.2	39.4	28.3 to 53.8	0.5	0.3 to 0.6
	Both	5 to 9	3.2	2.8 to 3.6	25.6	19.0 to 35.3	0.3	0.2 to 0.4
	Both	10 to 14	6.4	5.7 to 7.2	38.7	29.5 to 53.3	0.5	0.4 to 0.7
	Both	15 to 19	10.4	9.2 to 11.7	103.1	83.0 to 128.2	1.5	1.2 to 1.8
Oceania	Male	1 to 4	1.3	1.0 to 1.8	122.8	50.6 to 265.4	1.5	0.6 to 3.1

	Male	5 to 9	4	3.0 to 5.2	81.3	45.0 to 138.5	1	0.6 to 1.7
	Male	10 to 14	7.9	6.0 to 10.1	48.3	29.8 to 74.1	0.6	0.4 to 1.0
	Male	15 to 19	12.7	9.6 to 16.1	203.6	127.4 to 325.8	2.9	1.8 to 4.7
	Female	1 to 4	0.8	0.5 to 1.1	60.5	27.2 to 123.0	0.7	0.3 to 1.5
	Female	5 to 9	2.4	1.7 to 3.1	43.9	24.0 to 77.1	0.5	0.3 to 1.0
	Female	10 to 14	4.9	3.6 to 6.3	79.9	47.4 to 132.8	1.1	0.6 to 1.8
	Female	15 to 19	8.1	6.0 to 10.4	43	26.8 to 68.7	0.6	0.4 to 1.0
	Both	1 to 4	1.1	0.8 to 1.4	92.7	43.7 to 186.9	1.1	0.5 to 2.2
	Both	5 to 9	3.2	2.6 to 4.0	63.3	38.5 to 102.8	0.8	0.5 to 1.3
	Both	10 to 14	6.4	5.2 to 7.8	63.5	42.5 to 92.2	0.8	0.6 to 1.2
	Both	15 to 19	10.4	8.4 to 12.6	125.8	81.6 to 190.6	1.8	1.2 to 2.7
Central Asia	Male	1 to 4	0.7	0.6 to 0.8	20	13.6 to 28.2	0.2	0.2 to 0.3
	Male	5 to 9	2.2	1.8 to 2.6	20.9	15.3 to 28.0	0.3	0.2 to 0.3
	Male	10 to 14	4.5	3.7 to 5.3	20.1	15.0 to 26.9	0.3	0.2 to 0.4
	Male	15 to 19	7	5.7 to 8.4	76.5	55.9 to 102.5	1.1	0.8 to 1.5
	Female	1 to 4	0.5	0.4 to 0.6	18.1	9.5 to 26.3	0.2	0.1 to 0.3
	Female	5 to 9	1.4	1.2 to 1.7	19.5	12.0 to 25.5	0.2	0.1 to 0.3
	Female	10 to 14	2.7	2.2 to 3.2	23	16.0 to 32.4	0.3	0.2 to 0.4
	Female	15 to 19	4.1	3.3 to 5.1	46.4	32.5 to 62.5	0.7	0.5 to 0.9
	Both	1 to 4	0.6	0.5 to 0.7	19.1	14.0 to 25.0	0.2	0.2 to 0.3
	Both	5 to 9	1.8	1.6 to 2.1	20.2	15.9 to 24.8	0.3	0.2 to 0.3
	Both	10 to 14	3.6	3.2 to 4.1	21.5	17.3 to 26.8	0.3	0.2 to 0.4
	Both	15 to 19	5.6	4.9 to 6.4	61.8	49.0 to 76.9	0.9	0.7 to 1.1
Central Europe	Male	1 to 4	0.7	0.5 to 0.8	19.1	12.0 to 30.6	0.2	0.1 to 0.4
	Male	5 to 9	2.1	1.7 to 2.7	13.4	9.1 to 18.2	0.2	0.1 to 0.2
	Male	10 to 14	4.2	3.4 to 5.0	22.8	16.9 to 29.1	0.3	0.2 to 0.4
	Male	15 to 19	6.4	5.2 to 7.8	47.2	34.7 to 59.1	0.7	0.5 to 0.8
	Female	1 to 4	0.5	0.4 to 0.6	17.4	10.8 to 28.9	0.2	0.1 to 0.3

	Female	5 to 9	1.5	1.2 to 1.8	13.7	9.5 to 18.4	0.2	0.1 to 0.2
	Female	10 to 14	3	2.5 to 3.6	21.7	14.7 to 26.9	0.3	0.2 to 0.4
	Female	15 to 19	4.7	3.9 to 5.8	32	22.4 to 40.3	0.4	0.3 to 0.6
	Both	1 to 4	0.6	0.5 to 0.7	18.3	12.3 to 27.8	0.2	0.1 to 0.3
	Both	5 to 9	1.8	1.6 to 2.2	13.6	10.2 to 17.7	0.2	0.1 to 0.2
	Both	10 to 14	3.6	3.1 to 4.1	22.3	17.1 to 26.7	0.3	0.2 to 0.4
	Both	15 to 19	5.6	4.8 to 6.5	39.7	30.8 to 47.2	0.6	0.4 to 0.7
Eastern Europe	Male	1 to 4	0.4	0.2 to 0.7	3.7	2.0 to 5.7	0	0.0 to 0.1
	Male	5 to 9	1.3	0.7 to 2.1	5.2	2.9 to 7.8	0.1	0.0 to 0.1
	Male	10 to 14	2.6	1.6 to 3.9	9.9	6.3 to 14.0	0.1	0.1 to 0.2
	Male	15 to 19	4	2.4 to 6.0	40.4	27.2 to 56.7	0.6	0.4 to 0.8
	Female	1 to 4	0.3	0.2 to 0.6	4.7	1.6 to 7.4	0.1	0.0 to 0.1
	Female	5 to 9	1	0.5 to 1.9	7.2	4.5 to 11.9	0.1	0.1 to 0.1
	Female	10 to 14	1.9	1.2 to 3.1	12.4	6.4 to 17.1	0.2	0.1 to 0.2
	Female	15 to 19	2.9	1.8 to 4.9	36.2	19.1 to 49.5	0.5	0.3 to 0.7
	Both	1 to 4	0.4	0.2 to 0.6	4.2	2.3 to 6.1	0	0.0 to 0.1
	Both	5 to 9	1.2	0.7 to 1.8	6.2	4.3 to 8.3	0.1	0.1 to 0.1
	Both	10 to 14	2.3	1.6 to 3.1	11.1	7.7 to 14.1	0.1	0.1 to 0.2
	Both	15 to 19	3.4	2.4 to 4.9	38.3	27.8 to 48.7	0.5	0.4 to 0.7
High-income Asia Pacific	Male	1 to 4	1.8	1.3 to 2.5	10.5	6.5 to 15.8	0.1	0.1 to 0.2
	Male	5 to 9	5.4	3.9 to 6.9	7.3	4.8 to 9.8	0.1	0.1 to 0.1
	Male	10 to 14	10.5	7.9 to 13.3	14.1	9.5 to 18.2	0.2	0.1 to 0.2
	Male	15 to 19	17	12.7 to 22.0	22.5	16.0 to 30.1	0.3	0.2 to 0.4
	Female	1 to 4	1	0.7 to 1.4	7.9	5.2 to 13.8	0.1	0.1 to 0.2
	Female	5 to 9	3	2.2 to 3.9	9.4	6.1 to 12.9	0.1	0.1 to 0.2
	Female	10 to 14	6.1	4.7 to 7.8	10.4	7.7 to 15.4	0.1	0.1 to 0.2
	Female	15 to 19	10.2	7.6 to 13.0	17.2	12.7 to 26.9	0.2	0.2 to 0.4

	Both	1 to 4	1.4	1.1 to 1.8	9.3	6.4 to 13.4	0.1	0.1 to 0.2
	Both	5 to 9	4.2	3.3 to 5.1	8.3	6.2 to 10.4	0.1	0.1 to 0.1
	Both	10 to 14	8.4	6.8 to 10.0	12.3	9.7 to 15.1	0.1	0.1 to 0.2
	Both	15 to 19	13.7	11.1 to 16.6	20	15.8 to 25.0	0.3	0.2 to 0.3
Australasia	Male	1 to 4	1.4	0.9 to 2.0	2.6	1.5 to 4.1	0	0.0 to 0.0
	Male	5 to 9	4.1	2.8 to 5.5	2.5	1.7 to 3.6	0	0.0 to 0.0
	Male	10 to 14	8.1	5.8 to 10.4	5.8	4.0 to 7.9	0.1	0.0 to 0.1
	Male	15 to 19	13.4	9.3 to 17.6	14.2	9.0 to 19.7	0.2	0.1 to 0.2
	Female	1 to 4	0.9	0.6 to 1.3	2.6	1.5 to 3.7	0	0.0 to 0.0
	Female	5 to 9	2.6	1.8 to 3.7	3	2.0 to 4.4	0	0.0 to 0.1
	Female	10 to 14	5.2	3.6 to 7.0	5.8	3.9 to 7.8	0.1	0.0 to 0.1
	Female	15 to 19	8.5	5.9 to 11.8	12.9	7.6 to 17.5	0.2	0.1 to 0.2
	Both	1 to 4	1.1	0.9 to 1.5	2.6	1.7 to 3.7	0	0.0 to 0.0
	Both	5 to 9	3.4	2.6 to 4.3	2.8	2.1 to 3.6	0	0.0 to 0.0
	Both	10 to 14	6.7	5.3 to 8.3	5.8	4.3 to 7.4	0.1	0.0 to 0.1
	Both	15 to 19	11.1	8.6 to 13.6	13.5	9.4 to 17.2	0.2	0.1 to 0.2
Western Europe	Male	1 to 4	1.2	1.0 to 1.4	6.9	4.3 to 9.6	0.1	0.0 to 0.1
	Male	5 to 9	3.3	2.8 to 3.8	5.7	4.1 to 7.5	0.1	0.0 to 0.1
	Male	10 to 14	6.1	5.3 to 7.0	11.3	8.5 to 14.2	0.1	0.1 to 0.2
	Male	15 to 19	9.6	8.5 to 11.0	20.3	15.2 to 26.0	0.3	0.2 to 0.4
	Female	1 to 4	0.9	0.8 to 1.1	6.6	4.8 to 9.7	0.1	0.1 to 0.1
	Female	5 to 9	2.6	2.2 to 3.0	5.9	4.5 to 9.5	0.1	0.1 to 0.1
	Female	10 to 14	4.9	4.2 to 5.7	11.3	8.4 to 14.8	0.1	0.1 to 0.2
	Female	15 to 19	7.9	6.7 to 9.2	17.1	12.4 to 23.0	0.2	0.2 to 0.3
	Both	1 to 4	1	0.9 to 1.2	6.7	5.0 to 8.8	0.1	0.1 to 0.1
	Both	5 to 9	2.9	2.6 to 3.3	5.8	4.6 to 7.5	0.1	0.1 to 0.1
	Both	10 to 14	5.5	5.0 to 6.2	11.3	9.1 to 13.3	0.1	0.1 to 0.2

	Both	15 to 19	8.8	7.9 to 9.7	18.8	15.1 to 22.6	0.3	0.2 to 0.3
Southern Latin America	Male	1 to 4	1.3	1.0 to 1.8	25.8	14.1 to 43.1	0.3	0.2 to 0.5
	Male	5 to 9	3.8	2.9 to 4.8	22.3	14.6 to 31.5	0.3	0.2 to 0.4
	Male	10 to 14	7.2	5.6 to 8.9	34.3	24.3 to 47.0	0.4	0.3 to 0.6
	Male	15 to 19	11.4	8.7 to 14.2	79.1	54.8 to 112.0	1.1	0.8 to 1.6
	Female	1 to 4	0.6	0.4 to 0.8	15.3	8.8 to 26.4	0.2	0.1 to 0.3
	Female	5 to 9	1.6	1.1 to 2.3	16.7	11.1 to 24.9	0.2	0.1 to 0.3
	Female	10 to 14	3.1	2.3 to 4.2	28.4	20.5 to 40.6	0.4	0.3 to 0.5
	Female	15 to 19	5	3.6 to 6.6	49.8	33.4 to 68.1	0.7	0.5 to 1.0
	Both	1 to 4	1	0.7 to 1.2	20.6	12.4 to 32.3	0.2	0.1 to 0.4
	Both	5 to 9	2.7	2.1 to 3.3	19.6	14.0 to 26.4	0.2	0.2 to 0.3
	Both	10 to 14	5.2	4.2 to 6.2	31.4	24.3 to 40.2	0.4	0.3 to 0.5
	Both	15 to 19	8.2	6.7 to 9.9	64.7	48.7 to 82.7	0.9	0.7 to 1.2
High-income North America	Male	1 to 4	2.7	1.8 to 4.0	18.1	9.5 to 30.2	0.2	0.1 to 0.4
	Male	5 to 9	9.4	5.7 to 12.9	11.1	7.4 to 15.7	0.1	0.1 to 0.2
	Male	10 to 14	20.4	14.1 to 27.6	16.9	11.5 to 23.3	0.2	0.1 to 0.3
	Male	15 to 19	35.4	24.8 to 48.0	28.8	19.4 to 40.8	0.3	0.2 to 0.5
	Female	1 to 4	1.9	1.0 to 3.0	16.7	8.1 to 25.6	0.2	0.1 to 0.3
	Female	5 to 9	7.6	4.8 to 11.3	10	6.1 to 14.0	0.1	0.1 to 0.2
	Female	10 to 14	16.8	11.8 to 23.1	14.2	9.2 to 18.8	0.2	0.1 to 0.2
	Female	15 to 19	28.8	18.9 to 41.4	22.6	13.5 to 30.8	0.3	0.1 to 0.4
	Both	1 to 4	2.3	1.7 to 3.1	17.4	9.8 to 25.8	0.2	0.1 to 0.3
	Both	5 to 9	8.5	6.4 to 11.0	10.6	7.6 to 14.0	0.1	0.1 to 0.2
	Both	10 to 14	18.7	14.9 to 23.5	15.6	11.6 to 19.7	0.2	0.1 to 0.2

	Both	15 to 19	32.2	24.7 to 40.8	25.8	18.6 to 33.0	0.3	0.2 to 0.4
Caribbean	Male	1 to 4	0.8	0.6 to 1.0	252.5	121.2 to 459.1	3	1.4 to 5.4
	Male	5 to 9	2.5	2.0 to 3.1	89.4	52.8 to 139.5	1.1	0.7 to 1.8
	Male	10 to 14	4.8	4.0 to 6.0	80.6	54.3 to 117.5	1.1	0.7 to 1.6
	Male	15 to 19	7.6	6.3 to 9.5	81.1	53.9 to 117.4	1.2	0.8 to 1.7
	Female	1 to 4	0.6	0.5 to 0.8	237.8	125.8 to 419.6	2.8	1.5 to 5.0
	Female	5 to 9	1.9	1.5 to 2.4	82.6	50.3 to 128.3	1	0.6 to 1.6
	Female	10 to 14	3.6	2.9 to 4.3	69.3	44.3 to 100.7	0.9	0.6 to 1.3
	Female	15 to 19	5.5	4.4 to 6.6	79.4	51.4 to 117.2	1.1	0.7 to 1.7
	Both	1 to 4	0.7	0.6 to 0.9	245.3	148.7 to 379.9	2.9	1.8 to 4.5
	Both	5 to 9	2.2	1.9 to 2.6	86.1	59.4 to 120.1	1.1	0.7 to 1.5
	Both	10 to 14	4.2	3.7 to 5.0	75	56.8 to 98.9	1	0.8 to 1.3
	Both	15 to 19	6.6	5.7 to 7.7	80.3	59.4 to 104.9	1.1	0.8 to 1.5
Andean Latin America	Male	1 to 4	0.8	0.6 to 1.1	74.2	41.1 to 130.6	0.9	0.5 to 1.5
	Male	5 to 9	2.3	1.7 to 3.2	47.2	32.0 to 67.8	0.6	0.4 to 0.9
	Male	10 to 14	4.2	3.3 to 5.4	80.8	58.2 to 108.2	1.1	0.8 to 1.5
	Male	15 to 19	6.4	4.8 to 8.1	129.6	89.4 to 185.3	1.9	1.3 to 2.7
	Female	1 to 4	0.6	0.4 to 0.7	55	30.6 to 107.4	0.6	0.4 to 1.3
	Female	5 to 9	1.5	1.2 to 2.0	33.9	19.4 to 47.0	0.4	0.2 to 0.6
	Female	10 to 14	2.7	2.1 to 3.4	60.7	38.3 to 80.7	0.8	0.5 to 1.1
	Female	15 to 19	3.8	2.9 to 5.0	92.6	51.9 to 122.8	1.3	0.7 to 1.8
	Both	1 to 4	0.7	0.5 to 0.9	64.8	41.8 to 102.8	0.8	0.5 to 1.2
	Both	5 to 9	1.9	1.6 to 2.4	40.7	28.7 to 52.0	0.5	0.4 to 0.7
	Both	10 to 14	3.5	2.9 to 4.2	70.9	56.1 to 87.4	0.9	0.7 to 1.2
	Both	15 to 19	5.1	4.2 to 6.2	111.4	83.1 to 143.6	1.6	1.2 to 2.1
Central Latin America	Male	1 to 4	1.1	1.0 to 1.2	46.6	32.7 to 62.8	0.5	0.4 to 0.7
	Male	5 to 9	3.3	3.1 to 3.6	26.2	19.5 to 33.2	0.3	0.2 to 0.4

	Male	10 to 14	6.6	6.1 to 7.2	47.4	37.7 to 58.9	0.6	0.5 to 0.8
	Male	15 to 19	10.5	9.5 to 11.4	83.4	64.4 to 109.5	1.2	0.9 to 1.6
	Female	1 to 4	0.7	0.6 to 0.8	39.6	22.9 to 52.8	0.5	0.3 to 0.6
	Female	5 to 9	2.2	2.0 to 2.4	21.9	14.6 to 27.6	0.3	0.2 to 0.3
	Female	10 to 14	4.4	4.1 to 4.8	39	28.6 to 49.0	0.5	0.4 to 0.6
	Female	15 to 19	7.1	6.4 to 7.7	65.1	47.4 to 88.0	0.9	0.7 to 1.3
	Both	1 to 4	0.9	0.8 to 1.0	43.2	32.2 to 54.0	0.5	0.4 to 0.6
	Both	5 to 9	2.8	2.6 to 3.0	24.1	19.1 to 28.8	0.3	0.2 to 0.4
	Both	10 to 14	5.5	5.2 to 5.9	43.3	36.5 to 51.0	0.6	0.5 to 0.7
	Both	15 to 19	8.8	8.3 to 9.4	74.4	61.9 to 89.7	1.1	0.9 to 1.3
Tropical Latin America	Male	1 to 4	0.6	0.4 to 1.0	28.5	13.2 to 51.3	0.3	0.2 to 0.6
	Male	5 to 9	1.9	1.2 to 3.0	28.6	20.5 to 39.2	0.4	0.3 to 0.5
	Male	10 to 14	3.7	2.4 to 5.5	47.3	32.8 to 63.5	0.6	0.4 to 0.9
	Male	15 to 19	5.7	3.4 to 8.8	84.9	55.4 to 124.6	1.2	0.8 to 1.8
	Female	1 to 4	0.5	0.3 to 0.8	20.2	10.5 to 33.6	0.2	0.1 to 0.4
	Female	5 to 9	1.6	1.0 to 2.3	23.9	17.3 to 35.4	0.3	0.2 to 0.4
	Female	10 to 14	3	2.0 to 4.3	42.4	30.6 to 64.2	0.6	0.4 to 0.9
	Female	15 to 19	4.6	2.8 to 6.9	60	41.2 to 96.1	0.9	0.6 to 1.4
	Both	1 to 4	0.6	0.4 to 0.8	24.5	14.4 to 38.8	0.3	0.2 to 0.5
	Both	5 to 9	1.8	1.3 to 2.4	26.3	20.6 to 34.1	0.3	0.3 to 0.4
	Both	10 to 14	3.4	2.5 to 4.4	44.9	34.8 to 58.2	0.6	0.5 to 0.8
	Both	15 to 19	5.2	3.6 to 7.1	72.7	54.3 to 97.8	1	0.8 to 1.4
North Africa and Middle East	Male	1 to 4	0.1	0.1 to 0.2	227.8	160.8 to 315.6	2.7	1.9 to 3.7
	Male	5 to 9	0.4	0.3 to 0.5	158.8	128.4 to 193.5	2	1.6 to 2.4
	Male	10 to 14	0.8	0.6 to 0.9	171.4	142.8 to 205.7	2.3	1.9 to 2.8
	Male	15 to 19	1.3	1.0 to 1.6	279.2	221.5 to 343.0	4	3.2 to 5.0
	Female	1 to 4	0.1	0.1 to 0.1	195.2	141.1 to 274.2	2.3	1.7 to 3.2
	Female	5 to 9	0.3	0.2 to 0.4	113.7	90.0 to 156.5	1.4	1.1 to 2.0

	Female	10 to 14	0.7	0.5 to 0.8	107.8	85.6 to 145.7	1.4	1.2 to 2.0
	Female	15 to 19	1.1	0.8 to 1.3	135.2	100.0 to 175.7	1.9	1.4 to 2.5
	Both	1 to 4	0.1	0.1 to 0.1	211.9	163.8 to 273.7	2.5	1.9 to 3.2
	Both	5 to 9	0.4	0.3 to 0.4	136.7	113.3 to 162.4	1.7	1.4 to 2.0
	Both	10 to 14	0.7	0.6 to 0.8	140.4	120.7 to 167.3	1.9	1.6 to 2.3
	Both	15 to 19	1.2	1.0 to 1.4	209.1	175.9 to 244.5	3	2.5 to 3.5
South Asia	Male	1 to 4	1	0.7 to 1.4	61.2	31.9 to 108.4	0.7	0.4 to 1.3
	Male	5 to 9	3.1	2.2 to 4.0	46.2	27.1 to 76.0	0.6	0.3 to 1.0
	Male	10 to 14	6	4.4 to 7.8	94.4	61.7 to 145.0	1.3	0.8 to 1.9
	Male	15 to 19	9.2	6.6 to 12.5	146.1	94.0 to 221.1	2.1	1.3 to 3.2
	Female	1 to 4	0.7	0.5 to 0.9	46.8	20.7 to 113.6	0.6	0.2 to 1.3
	Female	5 to 9	2.1	1.5 to 2.8	69.5	37.7 to 140.6	0.9	0.5 to 1.8
	Female	10 to 14	4.3	3.3 to 5.8	75.1	42.9 to 131.4	1	0.6 to 1.8
	Female	15 to 19	7.2	5.3 to 9.5	108.7	63.4 to 174.6	1.6	0.9 to 2.5
	Both	1 to 4	0.9	0.7 to 1.1	54.4	31.0 to 96.1	0.6	0.4 to 1.1
	Both	5 to 9	2.6	2.1 to 3.2	57.3	36.8 to 94.5	0.7	0.5 to 1.2
	Both	10 to 14	5.2	4.2 to 6.2	85.2	59.6 to 121.8	1.1	0.8 to 1.6
	Both	15 to 19	8.2	6.6 to 10.1	128.2	92.5 to 174.7	1.8	1.3 to 2.5
Central Sub-Saharan Africa	Male	1 to 4	0.5	0.4 to 0.8	308.2	125.1 to 646.2	3.6	1.5 to 7.7
	Male	5 to 9	1.6	1.1 to 2.2	81.6	41.2 to 145.4	1	0.5 to 1.8
	Male	10 to 14	3	2.2 to 4.0	81.8	45.6 to 129.2	1.1	0.6 to 1.7
	Male	15 to 19	4.5	3.2 to 6.3	140.5	85.9 to 206.3	2	1.2 to 3.0
	Female	1 to 4	0.3	0.2 to 0.4	258.1	76.1 to 640.9	3.1	0.9 to 7.6
	Female	5 to 9	0.9	0.5 to 1.3	104.1	51.8 to 191.2	1.3	0.7 to 2.4
	Female	10 to 14	1.7	1.2 to 2.5	92.3	52.5 to 154.4	1.2	0.7 to 2.1
	Female	15 to 19	2.7	1.8 to 3.7	132.5	79.6 to 212.6	1.9	1.1 to 3.1
	Both	1 to 4	0.4	0.3 to 0.5	283.4	146.2 to 532.2	3.4	1.7 to 6.3
	Both	5 to 9	1.2	0.9 to 1.6	92.8	54.8 to 153.3	1.2	0.7 to 1.9

	Both	10 to 14	2.4	1.9 to 3.0	87.1	58.0 to 125.5	1.2	0.8 to 1.7
	Both	15 to 19	3.6	2.8 to 4.6	136.5	96.7 to 186.2	2	1.4 to 2.7
Eastern Sub-Saharan Africa	Male	1 to 4	0.6	0.5 to 0.8	17.1	9.6 to 28.6	0.2	0.1 to 0.3
	Male	5 to 9	1.8	1.5 to 2.2	15.4	9.5 to 22.5	0.2	0.1 to 0.3
	Male	10 to 14	3.1	2.6 to 3.6	21.6	14.5 to 30.4	0.3	0.2 to 0.4
	Male	15 to 19	4	3.4 to 4.8	82.9	58.1 to 111.5	1.2	0.8 to 1.6
	Female	1 to 4	0.3	0.2 to 0.3	13.3	6.4 to 29.4	0.2	0.1 to 0.3
	Female	5 to 9	0.8	0.7 to 1.0	10	6.2 to 17.6	0.1	0.1 to 0.2
	Female	10 to 14	1.5	1.3 to 1.8	12	7.5 to 19.6	0.2	0.1 to 0.3
	Female	15 to 19	2.2	1.9 to 2.6	38.2	22.0 to 56.8	0.5	0.3 to 0.8
	Both	1 to 4	0.5	0.4 to 0.5	15.2	9.5 to 24.3	0.2	0.1 to 0.3
	Both	5 to 9	1.3	1.2 to 1.5	12.7	8.9 to 17.2	0.2	0.1 to 0.2
	Both	10 to 14	2.3	2.0 to 2.6	16.8	12.0 to 22.2	0.2	0.2 to 0.3
	Both	15 to 19	3.1	2.7 to 3.5	60.6	45.6 to 77.0	0.9	0.7 to 1.1
Southern Sub-Saharan Africa	Male	1 to 4	0.4	0.3 to 0.6	162	69.7 to 318.3	1.9	0.8 to 3.8
	Male	5 to 9	1.3	0.8 to 1.8	45.1	24.2 to 76.1	0.6	0.3 to 1.0
	Male	10 to 14	2.6	1.8 to 3.5	40.4	25.0 to 62.1	0.5	0.3 to 0.8
	Male	15 to 19	4.2	2.9 to 5.8	104.3	67.6 to 148.0	1.5	1.0 to 2.1
	Female	1 to 4	0.3	0.2 to 0.4	293.6	56.9 to 637.0	3.5	0.7 to 7.5
	Female	5 to 9	0.9	0.6 to 1.3	57.2	21.7 to 113.3	0.7	0.3 to 1.4
	Female	10 to 14	2	1.4 to 2.7	36.1	17.9 to 58.7	0.5	0.2 to 0.8
	Female	15 to 19	3.3	2.3 to 4.6	70.1	39.0 to 103.0	1	0.6 to 1.5
	Both	1 to 4	0.3	0.2 to 0.4	227.2	77.8 to 413.8	2.7	0.9 to 4.9
	Both	5 to 9	1.1	0.8 to 1.4	51.1	25.8 to 83.5	0.6	0.3 to 1.1
	Both	10 to 14	2.3	1.8 to 2.8	38.3	24.0 to 55.0	0.5	0.3 to 0.7
	Both	15 to 19	3.7	2.9 to 4.7	87.3	61.2 to 115.6	1.3	0.9 to 1.7
Western Sub-Saharan Africa	Male	1 to 4	0.5	0.4 to 0.7	166.1	71.7 to 307.4	2	0.8 to 3.6
	Male	5 to 9	1.5	1.1 to 2.0	68.7	38.2 to 114.5	0.9	0.5 to 1.4

	Male	10 to 14	2.9	2.3 to 3.7	58.1	34.5 to 86.7	0.8	0.5 to 1.2
	Male	15 to 19	4.5	3.6 to 5.7	66.3	44.7 to 93.5	0.9	0.6 to 1.3
	Female	1 to 4	0.3	0.2 to 0.4	140.4	60.3 to 417.5	1.7	0.7 to 4.9
	Female	5 to 9	0.8	0.6 to 1.1	42.8	24.8 to 82.3	0.5	0.3 to 1.0
	Female	10 to 14	1.7	1.3 to 2.3	45.5	29.5 to 83.2	0.6	0.4 to 1.1
	Female	15 to 19	2.9	2.3 to 3.8	68.1	44.6 to 99.8	1	0.6 to 1.4
	Both	1 to 4	0.4	0.3 to 0.5	153.5	85.7 to 292.4	1.8	1.0 to 3.5
	Both	5 to 9	1.2	0.9 to 1.5	56	35.9 to 84.2	0.7	0.5 to 1.1
	Both	10 to 14	2.3	2.0 to 2.8	51.9	36.7 to 72.2	0.7	0.5 to 1.0
	Both	15 to 19	3.7	3.1 to 4.4	67.2	50.3 to 88.2	1	0.7 to 1.3