

Processes of change in cognitive therapy for PTSD

Taking a closer look at how therapy works

MentalHealth
ResearchUK

Scholar's Day 2019



Milan Wiedemann



@milanwiedemann


milan.wiedemann@gmail.com

Prof. Anke Ehlers and Prof. David M. Clark

Oxford Centre for Anxiety Disorders and Trauma



A journey from theory to therapy and back




Three approaches to explore process of change

Sudden gains


Sample 1

Sample 2




Mediation

Latent change score modeling



Network analysis

Cross-lagged network models




Three approaches to explore process of change

Sudden gains


Sample 1

Sample 2




Mediation

Latent change score modeling



Network analysis

Cross-lagged network models




Three approaches to explore process of change

Sudden gains


Sample 1

Sample 2




Mediation

Latent change score modeling




Network analysis

Cross-lagged network models



Automating the identification of sudden gains



<https://doi.org/10.31234/osf.io/2wa84>


suddengains: An R package to identify sudden gains in longitudinal data

Milan Wiedemann^{1,2}, Graham R Thew^{1,3}, Richard Stott⁴, & Anke Ehlers^{1,2,4}


Abstract

Sudden gains are large and stable changes on an outcome variable between consecutive measurements, for example during a psychological intervention with multiple sessions. Researching these occurrences could help understand individual change processes in longitudinal data. Tang and DeRubeis (1999) suggested three criteria to define sudden gains in psychological interventions. However, identifying sudden gains based on these criteria can be time consuming and prone to errors if not fully automated. Further, methodological decisions such as how missing data, or multiple gains, are handled vary across studies and are reported with different levels of detail. These problems limit the comparability of individual studies and make it hard to understand or replicate the exact methods used. The R package *suddengains* provides a set of tools to facilitate sudden gains studies by automating their identification, adjusting for missing data, and providing descriptive statistics as well as exportable data files for further analysis.

Keywords: sudden gains, sudden losses, R, mechanisms of change, processes of change, psychological therapy, longitudinal analysis, single case designs



* Definition based on Tang & DeRubeis (1999). Identification using the  package *suddengains*.

Cognitive processes associated with sudden gains





* Based on Ehlers and Clark (2000) and Ehlers et al. (2005). Description of Sample 1 in Ehlers et al. (2013).

Cognitive processes associated with sudden gains



* Based on Ehlers and Clark (2000) and Ehlers et al. (2005). Description of Sample 1 in Ehlers et al. (2013).

Cognitive processes associated with sudden gains



* Based on Ehlers and Clark (2000) and Ehlers et al. (2005). Description of Sample 1 in Ehlers et al. (2013).


Research Questions

How do cognitive processes^{*1} and comorbid symptoms^{*2} change **before, during, and after** sudden gains?

^{*1} Negative appraisals, memory characteristics.

^{*2} Depression and anxiety symptoms.

Baseline
Predictors of
sudden gains?




Differences in
treatment outcomes
at **end** and **follow up**?

Three approaches to explore process of change

Sudden gains


Sample 1

Sample 2




Mediation

Latent change score modeling




Network analysis

Cross-lagged network models




Treatment and samples




* Based on Ehlers and Clark (2000) and Ehlers et al. (2005).

Treatment and samples




* Based on Ehlers and Clark (2000) and Ehlers et al. (2005).

Research question: Mediation of clinical improvement



* Drawn from Ehlers and Clark's (2000) cognitive model of PTSD.

Research question: Mediation of clinical improvement





* Drawn from Ehlers and Clark's (2000) cognitive model of PTSD.

Bivariate latent change score modeling

Negative appraisals

PTSD symptoms





* Models were specified using the  package *lcs*m and analysed using *lavaan* (Rosseel, 2012). For method see Grimm et al. (2012).

Bivariate latent change score modeling


Negative appraisals

PTSD symptoms



* Models were specified using the  package *lcs*m and analysed using *lavaan* (Rosseel, 2012). For method see Grimm et al. (2012).

Summary of my journey so far




Three approaches to explore process of change

Sudden gains


Sample 1

Sample 2




Mediation

Latent change score modeling



Network analysis

Cross-lagged network models



Acknowledgements



MentalHealth
ResearchUK



THE AUSTIN AND HOPE PILKINGTON TRUST



THANK YOU



Milan Wiedemann



@milanwiedemann

milan.wiedemann@gmail.com

Prof. Anke Ehlers and Prof. David M. Clark

Oxford Centre for Anxiety Disorders and Trauma

