

Supplementary material for Kilander C, et al. Systematic underreporting of the population-based incidence of pancreatic and biliary tract cancers. Acta Oncol 2014;53:822–9.

Supplementary Table I. Cumulative mortality from date of diagnosis and concordance between diagnosis of pancreatic cancer in the Swedish Cancer Register (CR) and the Swedish Patient Register (PR) and cause of death in the Causes of Death Register within each survival time strata.

Sub population	Time since diagnosis (in years)					
	Total	0–0.5	0.5–1	1–3	3–5	> 5
PR-not-CR (number)	10 119	7806	1366	835	70	42
Proportion dead (%)	100	77	14	8	< 1	< 1
Concordance (%)	80	85	81	68	35	16
CR-not-PR (%)	757	658	65	29	4	1
Proportion dead (%)	100	87	9	4	< 1	< 1
Concordance (%)	71	73	68	56	50	11
CR-and-PR (%)	15 260	10 212	2829	1915	216	88
Proportion dead (%)	100	67	18	13	1	< 1
Concordance (%)	96	96	97	96	88	73

Percentages are rounded to whole figures.

Supplementary Table II. Cumulative mortality from date of diagnosis and concordance between diagnosis of biliary tract cancer in the Swedish Cancer Register (CR) and the Swedish Patient Register (PR) and cause of death in the Causes of Death Register within each survival time strata.

Sub population	Time since diagnosis (in years)					
	Total	0–0.5	0.5–1	1–3	3–5	> 5
PR-not-CR (number)	3438	2700	437	259	31	11
Proportion dead (%)	100	78	13	8	1	< 1
Concordance (%)	59	65	55	39	25	10
CR-not-PR (%)	552	464	53	29	2	4
Proportion dead (%)	100	84	10	5	< 1	< 1
Concordance (%)	51	58	41	33	10	9
CR-and-PR (%)	5358	3341	938	853	149	77
Proportion dead (%)	100	62	18	16	3	1
Concordance (%)	84	89	84	79	71	38

Percentages are rounded to whole figures.