

Supporting Information

Enhancing Photovoltaic Performance of Copolymers Containing Thiophene Unit with D-A Conjugated Side Chain by Rational Molecular Design

Ping Shen,^{1,2} Haijun Bin,^{1,2} Lu Xiao,¹ Yongfang Li^{1,3*}

¹Beijing National Laboratory for Molecular Sciences, CAS Key Laboratory of Organic Solids, Institute of Chemistry, Chinese Academy of Sciences, Beijing 100190, China

²College of Chemistry and Key Laboratory of Environmentally Friendly Chemistry and Applications of Ministry of Education, Xiangtan University, Xiangtan 411105, China

³College of Chemistry, Chemical Engineering and Materials Science, Soochow University, Suzhou, Jiangsu 215123, China.

*Correspondence to: Y. Li (E-mail: liyf@iccas.ac.cn)




Figure S1. Normalized photoluminescence spectra of the copolymer **P(BDT-TBTF/DPP)** excited at different $\lambda_{s,max}$ in dilute CHCl₃.




Figure S2. Cyclic voltammograms of the copolymer films on platinum electrode in acetonitrile solution containing 0.1 mol/L Bu_4NPF_6 at a scan rate of 100 mV/s. The cyclic voltammogram of ferrocene was also put in the figure for the potential calibration.